

CHARACTERIZATION OF CYCLICALLY FULLY COMMUTATIVE ELEMENTS IN FINITE AND AFFINE COXETER GROUPS

MATHIAS PÉTRÉOLLE

ABSTRACT. An element of a Coxeter group W is fully commutative if any two of its reduced decompositions are related by a series of transpositions of adjacent commuting generators. An element of a Coxeter group W is cyclically fully commutative if any of its cyclic shifts remains fully commutative. These elements were studied by Boothby *et al.* In particular the authors enumerated cyclically fully commutative elements in all Coxeter groups having a finite number of them. In this work we characterize and enumerate cyclically fully commutative elements according to their Coxeter length in all finite or affine Coxeter groups by using a new operation on heaps, the cylindric transformation. In finite types, this refines the work of Boothby *et al.*, by adding a new parameter. In affine type, all the results are new. In particular, we prove that there is a finite number of cyclically fully commutative logarithmic elements in all affine Coxeter groups. We study afterwards the cyclically fully commutative involutions and prove that their number is finite in all Coxeter groups.

INTRODUCTION

Let W be a Coxeter group. An element $w \in W$ is said to be fully commutative (FC) if any reduced word for w can be obtained from any other one by only short braid relations. These elements were introduced and studied independently by Fan in [4], Graham in [6] and Stembridge in [11, 12, 13], where among other things, he classified the Coxeter groups with a finite number of fully commutative elements and enumerated them in each case. Fully commutative elements appear naturally in the context of (generalized) Temperley-Lieb algebras, as they index a linear basis of the latter. Recently, in [1], Biagioli, Jouhet and Nadeau refined Stembridge's enumeration by counting fully commutative elements according to their Coxeter length in any finite or affine Coxeter group.

In this paper, we focus on a certain subset of fully commutative elements, the cyclically fully commutative elements (which we denote CFC from now on). These are elements for which every cyclic shift of any reduced expression is a reduced expression of a FC element. They were introduced by Boothby *et al.* in [3], where they classified the Coxeter groups with a finite number of CFC elements (they showed that the latter are the groups with a finite number of FC elements) and enumerated them. The main goal of [3] is to establish necessary and sufficient conditions for a CFC element $w \in W$ to be logarithmic, i.e to satisfy $\ell(w^k) = k\ell(w)$ for any positive integer k . This is the first step towards studying a cyclic version of

Key words and phrases. Cyclically fully commutative elements, Coxeter groups, generating functions, affine Coxeter groups, heaps.

Matsumoto's theorem for CFC elements. Here we will focus on the combinatorics of CFC elements. We introduce a new operation on heaps, which will allow us to give a new characterization of CFC elements in all Coxeter groups (see Theorem 1.11). For finite or affine Coxeter groups, this characterization can be reformulated in terms of words, by using the work on FC elements from [1]. It allows us to enumerate the CFC elements by taking into account their Coxeter length. We will also prove that the number of CFC involutions is finite in all Coxeter groups.

This paper is organized as follows. We recall in Section 1 some definitions and properties on Coxeter groups, we introduce a new operation on heaps, the *cylindric transformation*, in two different ways, and we deduce a new characterization of CFC elements in terms of pattern-avoidance for cylindric transformed heaps (see Theorem 1.11). In Section 2, we use this characterization in order to obtain a complete classification (in terms of words) of CFC elements in the affine group $\tilde{A}_{n-1}$. We also deduce a classification of CFC elements in the group A_{n-1} , and use this to enumerate CFC elements in both groups, according to their Coxeter length. The same work is done for the groups $\tilde{C}_n$, B_n , D_{n+1} , $\tilde{B}_{n+1}$, and $\tilde{D}_{n+2}$ in Section 3. In Section 4, we will focus on CFC involutions. The main result is that there is a finite number of CFC involutions in all Coxeter groups. We also give a characterization of CFC involutions for all Coxeter groups and enumerate them in finite and affine types.

1. CYCLICALLY FULLY COMMUTATIVE ELEMENTS AND HEAPS

1.1. Cyclically fully commutative elements.

Let W be a Coxeter group with a finite generating set S and Coxeter matrix $M = (m_{st})_{s,t \in S}$. Recall (see [2]) that this notation means that the relations between generators are of the form $(st)^{m_{st}} = 1$ for $m_{st} \neq \infty$, M is a symmetric matrix with $m_{ss} = 1$ and $m_{st} \in \{2, 3, \dots\} \cup \{\infty\}$. The pair (W, S) is called a Coxeter system. We can write the relations as $sts \dots = tst \dots$, each side having length m_{st} ; these are usually called braid relations. When $m_{st} = 2$, this is a commutation relation or a short braid relation. We define the Coxeter diagram Γ associated to (W, S) as the graph with vertex set S , and for each pair (s, t) with $m_{st} \geq 3$, an edge between s and t labelled by m_{st} . If $m_{st} = 3$, we usually omit the labelling.

For $w \in W$, we define the Coxeter length $\ell(w)$ as the minimum length of any expression $\mathbf{w} = s_1 \dots s_n$ with $s_i \in S$. An expression is called reduced if it has a minimal length. The set of all reduced expressions of w is denoted by $R(w)$. A classical result in Coxeter group theory, known as Matsumoto's theorem, is that any expression in $R(w)$ can be obtained from any other one using only braid relations. An element w is said to be fully commutative if any expression in $R(w)$ can be obtained from any other one using only commutation relations. For clarity, we will write w for an element in W , and $\mathbf{w}$ for one of its words (or expressions).

Definition 1.1. ([3, Definition 3.4]) *Given an expression $\mathbf{w} = s_{i_1} \dots s_{i_n}$, we define the left (resp. right) cyclic shift of $\mathbf{w}$ as $s_{i_2} \dots s_{i_n} s_{i_1}$ (resp. $s_{i_n} s_{i_1} \dots s_{i_{n-1}}$). A cyclic shift of $\mathbf{w}$ is $s_{i_k} \dots s_{i_{k-2}} s_{i_{k-1}}$ for $k \in \mathbb{N}$. An element $w \in W$ is said to be cyclically fully commutative if every cyclic shift of any expression in $R(w)$ is a reduced expression for a fully commutative element.*

We denote the set of CFC elements of W by W^{CFC} .

1.2. Heaps and FC elements.

We follow Stembridge [11] in this section. Fix a word $\mathbf{w} = (s_{a_1}, \dots, s_{a_\ell})$ in S^* , the free monoid generated by S . We define a partial ordering of the indices $\{1, \dots, \ell\}$ by $i \prec j$ if $i < j$ and $m(s_{a_i}, s_{a_j}) \geq 3$ and extend by transitivity on a partial ordering. We denote by $H_{\mathbf{w}}$ this poset together with the “labeling” $i \mapsto s_{a_i}$: this is the *heap* of $\mathbf{w}$. We will often call the elements of $H_{\mathbf{w}}$ points. In the Hasse diagram of $H_{\mathbf{w}}$, elements with the same labels will be drawn in the same column. The size $|H|$ of a heap H is its cardinality. Given any subset $I \subset S$, we will note H_I the subposet induced by all elements of H with labels in I . In particular $H_s := H_{\{s\}}$ for $s \in S$ is the chain $H_s = s^{(1)} \prec s^{(2)} \prec \dots \prec s^{(k)}$ where $k = |H_s|$. We also denote by $|\mathbf{w}_s|$ the number $|H_s|$ (note that this also counts the number of occurrences of s in $\mathbf{w}$). In H , we say that there is a *chain covering relation* between i and j , denoted by $i \prec_c j$, if $i \prec j$ and one of the two following conditions is satisfied:

- $m_{s_{a_i} s_{a_j}} \geq 3$ and there is no element z with the same label as i or j such that $i \prec z \prec j$,
- $s_{a_i} = s_{a_j}$ and there is no element z such that $i \prec z \prec j$.

Note that the set of chain covering relations corresponds to the set of edges in the corresponding heap. In other words, this set is equal to the set of covering relations of all subposets $H_{\{s,t\}}$, with $m_{st} \geq 3$. In Figure 1, we fix a Coxeter diagram on the left, and we give two examples of words with the corresponding heaps.




FIGURE 1. Two different heaps.

If we consider heaps up to poset isomorphisms which preserve the labels, then heaps encode precisely *commutativity classes*, that is, if the word $\mathbf{w}'$ is obtainable from $\mathbf{w}$ by transposing commuting generators then there exists a poset isomorphism between $H_{\mathbf{w}}$ and $H_{\mathbf{w}'}$. In particular, if w is fully commutative, the heaps of the reduced words are all isomorphic, and thus we can define the heap of w , denoted H_w .

Let $\mathbf{w} = (s_{a_1}, \dots, s_{a_\ell})$ be a word. A *linear extension* of the poset $H_{\mathbf{w}}$ is a linear ordering π of $\{1, \dots, \ell\}$ such that $\pi(i) < \pi(j)$ implies $i \prec j$. Now let $\mathcal{L}(H_{\mathbf{w}})$ be the set of words $(s_{a_{\pi(1)}}, \dots, s_{a_{\pi(\ell)}})$ where π ranges over all linear extensions of $H_{\mathbf{w}}$.

Proposition 1.2 ([11], Proposition 1.2). *Let w be a fully commutative element. Then $\mathcal{L}(H_{\mathbf{w}})$ is equal to $\mathcal{R}(w)$ for some (equivalently every) $\mathbf{w} \in \mathcal{R}(w)$.*

We say that a chain $i_1 \prec \dots \prec i_m$ in a poset H is *convex* if the only elements u satisfying $i_1 \preceq u \preceq i_m$ are the elements i_j of the chain. The next result characterizes *FC heaps*, namely the heaps representing the commutativity classes of FC elements.

Proposition 1.3 ([11], Proposition 3.3). *A heap H is the heap of some FC element if and only if (a) there is no convex chain $i_1 \prec_c \dots \prec_c i_{m_{st}}$ in H such that $s_{i_1} = s_{i_3} = \dots = s$ and $s_{i_2} = s_{i_4} = \dots = t$ where $3 \leq m_{st} < \infty$, and (b) there is no chain covering relation $i \prec_c j$ in H such that $s_i = s_j$.*

1.3. Cylindric transformation of heaps and CFC elements. Now, we will focus on CFC elements. Before this, we need to define a new operation on heaps.

Definition 1.4. Let $H := H_{\mathbf{w}}$ be a heap of a word $\mathbf{w} = s_{a_1} \dots s_{a_\ell}$. The cylindric transformation H^c of H is the labelling $i \mapsto s_{a_i}$ and a relation on the indices $\{1, \dots, \ell\}$, made of the chain covering relations $\prec_c$ of H , together with some new relations defined as follows:

- for each generator s , consider the minimal point a and the maximal point b in the chain H_s (for the partial order $\prec$). If a is minimal and b is maximal in the poset H , we add a new relation $b \prec_c a$.
- for each pair of generators (s, t) such that $m_{s,t} \geq 3$, consider the minimal point a and the maximal point b in the chain $H_{\{s,t\}}$ (for the partial order $\prec$). If these points have different labellings (one has label s and the other has label t), we add a new relation $b \prec_c a$.

Example: for the simply laced linear Coxeter diagram with 7 generators, consider the word $\mathbf{w} = s_2 s_1 s_0 s_3 s_2 s_6 s_5 s_4 s_5 s_6 s_3$. Its heap and its cylindric transformation are represented in Figure 2.




FIGURE 2. A heap and its cylindric transformation.

Remark 1.5. H^c is not a poset, even if we extend the relation $\prec_c$ by transitivity. Indeed, each point in H^c would otherwise be in relation with itself. As H^c is not a poset, we cannot draw its Hasse diagram. We define the diagram of H^c as the Hasse diagram of H , in which we add oriented edges representing the new relations described in Definition 1.4.

Let us explain the name “cylindric transformation” that we decided to use. Consider the Coxeter system (W, S) corresponding to the linear Coxeter diagram Γ_n of Figure 3.




FIGURE 3. The linear Coxeter diagram.

In this case, when we draw the diagram of H^c , it should be seen as drawn on a cylinder instead of a plane as for Hasse diagrams of heaps (in our cylinder, each chain H_s for a generator s can be seen as a circle).

As for heaps, if H^c is a cylindric transformation of a heap, the size $|H^c|$ is its cardinality. Given any subset $I \subset S$, we will write H_I^c for the set of points in H^c with labels in I . Unlike in the definition of H_I , this definition does not include the relations $\prec_c$ between those points.

We will now give the construction of the cylindric transformation of a heap in terms of words.

Proposition 1.6. *If $\mathbf{w} = s_{a_1} \dots s_{a_k}$ is a word and H its heap, let $\mathbf{w}_1 = s_{a_1} \dots s_{a_p}$ be the shortest prefix of $\mathbf{w}$ such that all generators which appear in $\mathbf{w}$ also appear in $\mathbf{w}_1$. We write $\mathbf{w}' = \mathbf{w}\mathbf{w}_1 := s_{c_1} \dots s_{c_{k+p}}$ where we denote by $c_i := a_i$, $1 \leq i \leq k$ and $c_{k+i} := a_i$, $1 \leq i \leq p$. Consider the heap $H_{\mathbf{w}'}$ of the word $\mathbf{w}'$. In $H_{\mathbf{w}'}$ we keep only the points and the chain covering relations, we identify the points i and $k+i$ for $1 \leq i \leq p$, and we denote the resulting diagram by H' . Then we have $H' = H^c$.*

Remark 1.7. Note that H' define in Proposition 1.6 is not a poset, although $H_{\mathbf{w}'}$ is.




FIGURE 4. Illustration of Proposition 1.6 for $\mathbf{w} = s_1 s_0 s_1 s_3 s_2 s_6 s_5 s_4 s_6 s_5 s_3 s_0 s_1 s_0$. We have $\mathbf{w}_1 = s_1 s_0 s_1 s_3 s_2 s_6 s_5 s_4$.

Proof. We will first prove that the chain covering relations $\prec_c$ in H' are included in those of H^c . There are three types of chain covering relations in $H_{\mathbf{w}'}$:

- The ones between two points i and j such that $1 \leq i, j \leq k$. They correspond to chain covering relations which also appear in H , and are therefore present in H^c .
- The ones between two points i and j such that $k+1 \leq i, j \leq k+p$. After the identification between i (resp. j) and $i-k$ (resp. $j-k$), these relations also appear in H , and so in H^c .
- The ones between two points i and j such that $1 \leq i \leq k$ and $k+1 \leq j \leq p$. By construction of H' , the point j will be identified with the point $h := j-k$ which is minimal in $H_{s_{a_j}}$ (and even minimal in H if $s_{a_j} = s_{a_i}$). Likewise, the point i is maximal in $H_{s_{a_i}}$ (and even maximal in H if $s_{a_j} = s_{a_i}$). So, when we construct H^c , we will add a chain covering relation $i \prec_c h$.

Thus the chain covering relations of H' are included in those of H^c . The reverse inclusion can be proved by similar arguments. $\square$

Proposition 1.8. *The canonical mapping $H \mapsto H^c$ is not injective. More precisely, if the word $\mathbf{w}_2$ is commutation equivalent to some cyclic shift of the word $\mathbf{w}_1$, then $H_{\mathbf{w}_1}^c = H_{\mathbf{w}_2}^c$.*

Proof. For s a generator, $\mathbf{w}$ a word, define $H := H_{s\mathbf{w}}$ and $H' := H_{\mathbf{w}s}$. We will prove that the chain covering relations of H^c are included in those of H'^c . By applying this to the successive right cyclic shifts of $\mathbf{w}s$, we may conclude that $H^c = H'^c$,

which is sufficient to prove the proposition.

We denote by i (*resp.* i') the point of H (*resp.* H') corresponding to the first (*resp.* last) occurrence of s in $\mathbf{sw}$ (*resp.* $\mathbf{ws}$). There are three types of chain covering relations in H^c :

- The ones between two points different from i ; they are also present in H'^c .
- The ones between i and another point j . Two cases can occur: $i \prec_c j$ in H or not. In the first case, j must be minimal among all points with the same label. As i' is maximal in H' , we add a relation $i' \prec_c j$ in H'^c when we construct H'^c . The second case occurs when we add a relation between i and j during the construction of H^c . In this case, j is maximal in H among all points with the same label and so there is a relation $j \prec_c i'$ in H' . This relation stays naturally present in H'^c .
- The ones between i and itself. It is clear that i' is also in relation with itself in H'^c .

□

Now we characterize the CFC elements in terms of heaps.

Definition 1.9. Set H a heap and s, t two generators such that $m_{st} \geq 3$. Consider a chain $i_1 \prec_c \dots \prec_c i_m$ in H^c with $m \geq 3$, all elements i_k distinct and such that $s_{a_{i_1}} = s_{a_{i_3}} = \dots = s$ and $s_{a_{i_2}} = s_{a_{i_4}} = \dots = t$.

Such a chain is called *cylindric convex* if the only elements $u_1, \dots, u_d$, satisfying $i_1 \prec_c \dots \prec_c i_k \prec_c u_1 \prec_c \dots \prec_c u_d \prec_c i_m$ with all elements involved in this second chain distinct, are the elements i_j of the first chain.

Lemma 1.10. Assume there exists a convex chain $i_1 \prec_c \dots \prec_c i_m$ of length $m \geq 3$ in a heap H , such that all elements i_k are distinct and $s_{a_{i_1}} = s_{a_{i_3}} = \dots = s$ and $s_{a_{i_2}} = s_{a_{i_4}} = \dots = t$. Then this chain is also cylindric convex in the cylindric transformation H^c .

Proof. Let $i_1 \prec_c \dots \prec_c i_m$ be such a convex chain in H and assume there is a chain $i_1 \prec_c \dots \prec_c i_k \prec_c u_1 \prec_c \dots \prec_c u_d \prec_c i_m$ with all elements distinct in H^c , and $u_1 \neq i_{k+1}$. Two cases can occur:

- $i_k \neq i_{m-1}$: as i_k is not a maximal element in $H_{s_{a_{i_k}}}$, the relation $i_k \prec_c u_1$ holds in H . Consequently, the relation $i_k \prec_c u_1 \prec_c i_{k+2}$ holds in H . So $i_1 \prec_c \dots \prec_c i_m$ is not a convex chain in H , which is a contradiction,
- $i_k = i_{m-1}$: as i_m is not a minimal element in $H_{s_{a_{i_m}}}$, the relation $u_d \prec_c i_m$ holds in H . Consequently, the relation $i_{m-2} \prec_c u_d \prec_c i_m$ holds in H . So, as $u_d \neq i_{m-1}$, $i_1 \prec_c \dots \prec_c i_m$ is not a convex chain in H , which is a contradiction.

□

Theorem 1.11. A heap H is the heap of some word $\mathbf{w}$ corresponding to a CFC element w if and only if both the following conditions are satisfied:

- there is no cylindric convex chain $i_1 \prec_c \dots \prec_c i_{m_{st}}$ in the cylindric transformation H^c such that $s_{a_{i_1}} = s_{a_{i_3}} = \dots = s$ and $s_{a_{i_2}} = s_{a_{i_4}} = \dots = t$, where $3 \leq m_{st} < \infty$;
- there is no chain covering relation $i \prec_c j$ in the cylindric transformation H^c such that $s_{a_i} = s_{a_j}$.

Proof. Let $\mathbf{w}$ be a word and $H := H_{\mathbf{w}}$ its heap. Assume that $\mathbf{w}$ is a reduced word of a non CFC element. There exists by Definition 1.1 a cyclic shift of $\mathbf{w}$ which is commutation equivalent to $\mathbf{w}' = w_1 s s w_2$ or $\mathbf{w}' = w_1 \underbrace{st \cdots st}_{m_{st}} w_2$. Let H' be the heap

of $\mathbf{w}'$. By Proposition 1.3, in the first case, H' contains a chain covering relation $i \prec_c j$ such that $s_{a_i} = s_{a_j} = s$, and in the second case, H' contains a convex chain $i_1 \prec_c \cdots \prec_c i_{m_{st}}$ such that $s_{a_{i_1}} = s_{a_{i_3}} = \cdots = s$ and $s_{a_{i_2}} = s_{a_{i_4}} = \cdots = t$, where $3 \leq m_{st} < \infty$. By Lemma 1.10, these two cases give a chain covering relation and a cylindric convex chain in H'^c , respectively. But $\mathbf{w}'$ is commutation equivalent to a cyclic shift of $\mathbf{w}$, so $H^c = H'^c$, as seen in Proposition 1.8.

Conversely, if H^c contains the cylindric convex chain $i_1 \prec_c \cdots \prec_c i_{m_{st}}$, let $\mathbf{w}'$ be the cyclic shift of $\mathbf{w}$ beginning by s_{i_1} , and let H' be its heap. As all elements in the cylindric convex chain are distinct, H' contains the convex chain $i_1 \prec_c \cdots \prec_c i_{m_{st}}$. Therefore H' is not FC by Proposition 1.3 and H is not CFC. The same argument also holds if H^c contains a relation $i \prec_c j$ such that $s_{a_i} = s_{a_j}$, by letting this time $\mathbf{w}'$ be the cyclic shift of $\mathbf{w}$ beginning by s_{a_i} . $\square$

For example, this theorem ensures that the heap in Figure 2 does not correspond to a CFC element, as it contains a chain covering relation $i \prec_c j$ in the cylindric transformation H^c such that $s_{a_i} = s_{a_j} = s_6$. The example in Figure 5 corresponds to a CFC element, according to Theorem 1.11.




FIGURE 5. A CFC element in $\tilde{C}_6$.

Before ending this section, let us recall the set of alternating heaps which was defined in [1], and will be useful later.

Definition 1.12. In a Coxeter group with linear diagram Γ_n , a reduced word is alternating if for $i = 0, 1, \dots, n-1$, the occurrences of s_i alternate with those of s_{i+1} . A heap is called alternating if it is the heap of an alternating word. If the Coxeter group is $\tilde{A}_{n-1}$ (see Figure 6 for the Coxeter diagram), the diagram is not linear but we define the alternating word in the same way by setting $s_0 = s_n$.

Remark 1.13. The subset of alternating words defined as the set of words in which each generator occurs at most once is particular, in the sense that each element of this subset is a reduced expression for a CFC element (we can not use any braid relation of length at least 3). This fact justifies why this subset will be treated separately from other alternating words in the rest of the article.

2. CFC ELEMENTS IN TYPES $\tilde{A}$ AND A

In this section, we will give a characterization and the enumeration of CFC elements in both types A_{n-1} and $\tilde{A}_{n-1}$. The corresponding Coxeter diagrams are given in Figure 6.




FIGURE 6. Coxeter diagrams of types A_{n-1} and $\tilde{A}_{n-1}$.

The elements of A_{n-1}^{CFC} were enumerated in [3] by using recurrence relations. Our characterization in terms of heaps allows us to take into account the Coxeter length. Actually, we can compute the generating functions $W^{CFC}(q) = \sum_{w \in W^{CFC}} q^{\ell(w)}$ for $W = A_{n-1}$ and $\tilde{A}_{n-1}$. In particular, when $q = 1$ and $W = A_{n-1}$, we get back the enumeration from [3] (recall that the number of CFC elements in type $\tilde{A}_{n-1}$ is infinite). Our strategy is the following: first, we obtain a characterization of CFC elements in type $\tilde{A}_{n-1}$, we deduce from it a characterization of CFC elements in type A_{n-1} , then we derive the enumeration of CFC elements in type A_{n-1} and deduce from it the enumeration in type $\tilde{A}_{n-1}$.

2.1. Characterization in type $\tilde{A}_{n-1}$. Note that, in this type, the diagram of a cylindric transformation of a heap can be seen as drawn on a torus.

Theorem 2.1. *An element $w \in \tilde{A}_{n-1}$ is CFC if and only if one (equivalently, any) of its reduced expressions $\mathbf{w}$ verifies one of these conditions:*

- (a) *each generator occurs at most once in $\mathbf{w}$,*
- (b) *$\mathbf{w}$ is an alternating word and $|\mathbf{w}_{s_0}| = |\mathbf{w}_{s_1}| = \dots = |\mathbf{w}_{s_{n-1}}| \geq 2$.*

Proof. As said in Remark 1.13, if each generator occurs at most once in $\mathbf{w}$, then w is a CFC element. So let $\mathbf{w}$ be a reduced expression of a CFC element w having a generator occurring at least twice in $\mathbf{w}$. Recall that according to [1], $w \in \tilde{A}_{n-1}$ is fully commutative if and only if $\mathbf{w}$ is an alternating word. Let s_j be a generator which occurs at least twice in $\mathbf{w}$ and such that for all $k \in \{0, 1, \dots, n-1\}$, $|\mathbf{w}_{s_j}| \geq |\mathbf{w}_{s_k}|$. We will prove that $|\mathbf{w}_{s_j}| = |\mathbf{w}_{s_{j+1}}|$ where we set $s_n = s_0$, which is sufficient to show that each generator occurs the same number of times. As $\mathbf{w}$ is alternating, there are three possibilities:

- $|\mathbf{w}_{s_j}| = |\mathbf{w}_{s_{j+1}}| - 1$. It contradicts the maximality of $|\mathbf{w}_{s_j}|$.
- $|\mathbf{w}_{s_j}| = |\mathbf{w}_{s_{j+1}}| + 1$. By maximality of $|\mathbf{w}_{s_j}|$, there are two possibilities for $H_{\mathbf{w}}^c$: $|\mathbf{w}_{s_j}| = |\mathbf{w}_{s_{j-1}}|$ or $|\mathbf{w}_{s_j}| = |\mathbf{w}_{s_{j-1}}| + 1$. We obtain either a cylindric convex chain $x \prec_c y \prec_c z$ where x and z have label s_j and y has label s_{j-1} , or a chain covering relation between two indices p and q with label s_j , which allows us to conclude that w is not a CFC element by Theorem 1.11 (see the figure below, where we have circled the points x, y, z , and p, q respectively).




FIGURE 7. The two possible heaps.

- $|\mathbf{w}_{s_j}| = |\mathbf{w}_{s_{j+1}}|$ which is the needed condition.

Conversely, let $\mathbf{w}$ be an alternating word such that $|\mathbf{w}_{s_0}| = |\mathbf{w}_{s_1}| = \dots = |\mathbf{w}_{s_{n-1}}| \geq 2$. $H_{\mathbf{w}}^c$ can not contain a cylindric convex chain $x \prec_c y \prec_c z$ of length 3 such that $s_{a_x} = s_{a_z} = s_m$ and $s_{a_y} = s_{m+1}$ (resp. s_{m-1}): indeed, the required condition on $\mathbf{w}$ implies that there exists an index ℓ such that $x \prec_c \ell \prec_c z$ with $s_{a_\ell} = s_{m-1}$ (resp. s_{m+1}), which is a contradiction with the cylindric convexity of the chain. The same argument holds for chain covering relations involving indices with the same labellings. $\square$

2.2. Characterization and enumeration in type A_{n-1} . In this case, we will both characterize the CFC elements and compute the generating function

$$A^{CFC}(x) := \sum_{n \geq 1} A_{n-1}^{CFC}(q)x^n.$$

We begin with a lemma which is a consequence of Corollary 5.6 in [3]. Nevertheless, we will give a shorter proof using Theorem 2.1.

Lemma 2.2. *The CFC elements in type A_{n-1} are those having reduced expressions in which each generator occurs at most once.*

Proof. Let $\mathbf{w}$ be a reduced word of a CFC element in type A_{n-1} . By definition of the Coxeter diagram (cf Figure 6), it is a reduced word of a CFC element in type $\tilde{A}_{n-1}$ in which the generator s_0 does not occur. According to Theorem 2.1, the only such CFC elements in type $\tilde{A}_{n-1}$ are those in which each generator occurs at most once. Conversely, if all generators occur at most once in $\mathbf{w}$, Remark 1.13 ensures that $\mathbf{w}$ is a reduced expression of a CFC element. $\square$

Theorem 2.3. *We have $A_0^{CFC}(q) = 1$, $A_1^{CFC}(q) = 1 + q$ and for $n \geq 3$,*

$$A_{n-1}^{CFC}(q) = (2q + 1)A_{n-2}^{CFC}(q) - qA_{n-3}^{CFC}(q). \quad (1)$$

Equivalently, we have the generating function:

$$A^{CFC}(x) = x \frac{1 - qx}{1 - (2q + 1)x + qx^2}.$$

Proof. According to [1, Proposition 2.7], FC elements in A_{n-1} are in bijection with Motzkin type paths of length n , with starting and ending points at height 0, where the horizontal steps are labeled either L or R (and horizontal steps at height 0 are always labeled R). We recall the bijection, which is defined as follows: let w be a FC element in A_{n-1} , and H its heap. To each $s_i \in S$, we associate a point $P_i = (i, |H_{s_i}|)$. As w is alternating, three cases can occur:

- $|H_{s_i}| = |H_{s_{i+1}}| - 1$, corresponding to an ascending step,
- $|H_{s_i}| = |H_{s_{i+1}}| + 1$, corresponding to a descending step,
- $|H_{s_i}| = |H_{s_{i+1}}|$, corresponding to an horizontal step, labelled by R if s_i occurs before s_{i+1} in w , L otherwise.

According to Lemma 2.2, the restriction of this bijection to CFC elements is a bijection between CFC elements and the previous Motzkin paths, having length n , whose height does not exceed 1. By taking into account the first return to the x -axis, we obtain the following recurrence relation for $n \geq 3$:

$$A_{n-1}^{CFC}(q) = A_{n-2}^{CFC}(q) + \sum_{m=2}^n 2^{m-2} q^{m-1} A_{n-1-m}^{CFC}(q), \quad (2)$$

where we write $A_{-1}^{CFC}(q) = 1$ (which fits with $A_0^{CFC}(q) = 1$, $A_1^{CFC}(q) = 1 + q$ and the expected recurrence relation). Rewriting (2) with n replaced by $n - 1$ and combining with (2) allows us to eliminate the sum over m , and leads to the recurrence relation (1).

Classical techniques in generating function theory and the values $A_0^{CFC}(q) = 1$, $A_1^{CFC}(q) = 1 + q$ enable us to compute the desired generating function. $\square$

Notice that, as expected, if $q \rightarrow 1$, we find the odd-index Fibonacci numbers generating function of [3]. This q -analog was already known, see sequence A105306 in [10].

2.3. Enumeration in type $\tilde{A}_{n-1}$.

Proposition 2.4. *We have for $n \geq 3$*

$$\tilde{A}_{n-1}^{CFC}(q) = P_{n-1}(q) + \frac{2^n - 2}{1 - q^n} q^{2n}, \quad (3)$$

where $P_{n-1}(q)$ is a polynomial of degree n satisfying for $n \geq 4$

$$P_n(q) = (3q + 1)P_{n-1}(q) + (2q + 2q^2)P_{n-2}(q) - q^2 P_{n-3}(q),$$

with $P_1(q) = 1 + 2q + 2q^2$, $P_2(q) = 1 + 3q + 6q^2 + 6q^3$, and $P_3(q) = 1 + 4q + 10q^2 + 16q^3 + 14q^4$. Moreover, we can compute the generating function:

$$P(x) := \sum_{n=1}^{\infty} P_n(q)x^n = \frac{x(1 + 2q + 2q^2 - (2q + 2q^2)x + q^2 x^2)}{(1 - qx)(1 - (2q + 1)x + qx^2)}.$$

Therefore the coefficients of $\tilde{A}_{n-1}^{CFC}(q)$ are ultimately periodic of exact period n , and the periodicity starts at length n .

Proof. In the same way as for the finite type A_{n-1} (see [1]), FC elements in $\tilde{A}_{n-1}$ are in bijection with Motzkin type paths of length n satisfying the following conditions:

- the starting point $P_0 = (0, |H_{s_0}|)$ and the ending point $P_n = (n, |H_{s_n}| = |H_{s_0}|)$ have the same height,
- horizontal steps at height 0 are always labeled R,

- if the path contains only horizontal steps at height ≥ 1 , the steps must have not all the same labelling.

The construction of the path corresponding to a FC element is the same as in type A if we set $s_n = s_0$. In Theorem 2.1, the alternating CFC elements in which all generators occur at least twice and in the same number correspond to Motzkin type paths which have only horizontal steps. Therefore there are $2^n - 2$ such paths for all fixed starting height $h \geq 2$. It leads to the generating function

$$\sum_{h=2}^{+\infty} (2^n - 2)(q^n)^h,$$

which can be summed to obtain the second term of the right hand side of (3). In Theorem 2.1, the CFC elements which correspond to reduced expressions with at most one occurrence of each generator correspond to Motzkin type paths which stay at height ≤ 1 . We denote by $P_{n-1}(q)$ the generating function of such elements. Let i (*resp.* j) be the first (*resp.* last) return to the x-axis (see Figure 8 for an example).




FIGURE 8. Motzkin path corresponding to the element $s_{14}s_1s_0s_2s_6s_5s_7s_9s_{12}s_{11}$.

We get

$$\begin{aligned} P_{n-1}(q) = & (2^n - 2)q^n + nq^{n-1}2^{n-2} \\ & + \sum_{\substack{i=0 \\ i \neq 0 \text{ or } j \neq n-1}}^{n-2} \sum_{\substack{j=i+1 \\ j \neq n-1}}^{n-1} q^{n-1+i-j} 2^{n-2+i-j} A_{j-i-1}^{CFC}(q) + A_{n-2}^{CFC}(q), \end{aligned} \quad (4)$$

where the first term counts paths which stay at height 1, the second term counts paths such that $i = j$, and the last term counts paths such that $i = 0$ and $j = n - 1$. Rewriting (4) with $n - 1$ replaced by $n - 2$, combining with (4) and using the relation (2) in Theorem 2.3 allow us to eliminate the double sum, and leads to the following relation:

$$P_{n-1}(q) = 2qP_{n-2}(q) + 2q^n + A_{n-1}^{CFC}(q) - qA_{n-3}^{CFC}(q).$$

From this and (1), it is easy to derive the expected relation and the generating function P . As $P_{n-1}(q)$ is a polynomial in q , the periodicity is clear, and its beginning comes from the fact that the leading coefficient of $P_{n-1}(q)$ is $(2^n - 2)q^n$ and $[q^{n-1}]P_{n-1}(q) \neq 0$, where $[q^{n-1}]P_{n-1}(q)$ is the coefficient of q^{n-1} in $P_{n-1}(q)$. $\square$

The situation in type $\tilde{A}_{n-1}$ is very different from all other types that we study, as we will see later: this is the only case where the generating series P_n of CFC elements which correspond to reduced expression with at most one occurrence of each generator do not satisfy the recurrence relation $f_n(q) = (2q + 1)f_{n-1}(q) - qf_{n-2}(q)$.

3. TYPES $\tilde{C}$, $\tilde{B}$, $\tilde{D}$, B AND D .

There are three other infinite families of affine Coxeter groups, they correspond to types $\tilde{B}$, $\tilde{C}$, $\tilde{D}$. All these groups contain an infinite number of CFC elements. In any of these cases, using Theorem 1.11, we are able to characterize the CFC elements and to compute the generating function $W^{CFC}(q) = \sum_{w \in W^{CFC}} q^{\ell(w)}$. We also show that this generating function is always ultimately periodic. As a corollary, we deduce a characterization and the enumeration of CFC elements in finite types B and D .

3.1. Types $\tilde{C}_n$ and B_n . The Coxeter diagram of type $\tilde{C}_n$ is represented below.




FIGURE 9. Coxeter diagram of type $\tilde{C}_n$.

The situation is more complicated than in type $\tilde{A}_{n-1}$, but we obtain an explicit characterization of CFC elements.

Theorem 3.1. *An element $w \in \tilde{C}_n$ is CFC if and only if one (equivalently any) of its reduced expressions $\mathbf{w}$ verifies one of the three conditions:*

- (a) *each generator occurs at most once in $\mathbf{w}$,*
- (b) *$\mathbf{w}$ is an alternating word and $|\mathbf{w}_t| = |\mathbf{w}_{s_1}| = \dots = |\mathbf{w}_{s_{n-1}}| = |\mathbf{w}_u| \geq 2$,*
- (c) *$\mathbf{w}$ is a subword of the infinite periodic word $(ts_1s_2 \dots s_{n-1}us_{n-1} \dots s_2s_1)^\infty$, where $|\mathbf{w}_{s_1}| = \dots = |\mathbf{w}_{s_{n-1}}| \geq 2$ and $|\mathbf{w}_t| = |\mathbf{w}_u| = |\mathbf{w}_{s_1}|/2$ (i.e we have $\mathbf{w} = s_i s_{i+1} \dots s_{i-2} s_{i-1}$ or $s_i s_{i-1} \dots s_{i+2} s_{i+1}$, where $s_0 = t$ and $s_n = u$).*

Proof. Let w be a CFC element in $\tilde{C}_n$ and $\mathbf{w}$ be one of its reduced expressions. We denote by H the heap of $\mathbf{w}$ and H^c its cylindric transformation. In [1, Theorem 3.4], FC elements are classified in five families, the first corresponding to alternating elements. As before, we distinguish two cases for elements in this first family:

- each generator occurs at most once in $\mathbf{w}$. According to Remark 1.13, w is CFC as no long braid relations can be applied. These elements satisfy (a).
- $\mathbf{w}$ is an alternating word in which a generator occurs at least twice. In this case, the proof will essentially be the same as for alternating words in type $\tilde{A}_{n-1}$. We write $t = s_0$ and $u = s_n$. Let $\llbracket i_k, i_\ell \rrbracket$ be a maximal interval such that $|\mathbf{w}_{s_{i_k}}| = \dots = |\mathbf{w}_{s_{i_\ell}}| \geq 2$ and $\forall j \in \{0, 1, \dots, n\}$, $|\mathbf{w}_{s_j}| \leq |\mathbf{w}_{s_{i_\ell}}|$. Assume $i_\ell \leq n-1$, by maximality of $|\mathbf{w}_{s_{i_\ell}}|$ and the fact that $\mathbf{w}$ is alternating, we have $|\mathbf{w}_{s_{i_\ell}}| = |\mathbf{w}_{s_{i_\ell+1}}| + 1$ and there are two possibilities in H^c : $|H^c_{s_{i_\ell}}| = |H^c_{s_{i_\ell-1}}|$ or $|H^c_{s_{i_\ell}}| = |H^c_{s_{i_\ell-1}}| + 1$. We obtain either a cylindric convex chain $v \prec_c x \prec_c y \prec_c z$ of length 4 where v, y both have label s_{i_ℓ} and x, z both have label $s_{i_\ell-1}$, or a chain covering relation between two indices p and q with label s_{i_ℓ} , which allows us to conclude that w is not CFC by Theorem 1.11 (see figure below, in each case, we have circled the points v, x, y, z and q, r respectively).

So, i_ℓ has to be equal to n if w is CFC. The same argument for i_k will lead to $i_k = 0$, and so $|\mathbf{w}_{s_0}| = |\mathbf{w}_{s_1}| = \dots = |\mathbf{w}_{s_{n-1}}| \geq 2$. These elements satisfy (b).





FIGURE 10. The two possible heaps.

Next, we will show that among the four remaining families from [1], the CFC elements must satisfy (c). There are two cases:

- w belongs to the family called zigzag, i.e $\mathbf{w}$ is a subword of the infinite periodic word $(ts_1s_2 \dots s_{n-1}us_{n-1} \dots s_2s_1)^\infty$ with at least one generator which occurs more than three times (actually, this condition will not be used in this proof). Denote by s_i (*resp.* s_j) the first (*resp.* last) letter of $\mathbf{w}$. We assume that the second letter is s_{i-1} (the case where the second letter is s_{i+1} is symmetric and can be treated similarly).

If $s_j \notin \{s_{i-1}, s_{i+1}\}$, H^c contains a cylindric convex chain of length 3 involving points in $H^c_{\{s_{i-1}, s_i\}}$ or a chain covering relation between two points with label s_i .

If $s_j = s_{i-1}$, then the last but one letter in $\mathbf{w}$ is either s_i or s_{i-2} , so H^c takes one of the two forms in Figure 11.

FIGURE 11. The two possible heaps for $s_j = s_{i-1}$.

In the first case, H^c contains a cylindric convex chain involving points in $H_{\{s_{i-1}, s_i\}}^c$ of length 4, given by the two first and the two last letters of $\mathbf{w}$ (the corresponding points are circled in Figure 11, left). By Theorem 1.11, $\mathbf{w}$ is not CFC. In the second case, H^c contains a cylindric convex chain involving points in $H_{\{s_{i-1}, s_i\}}^c$ of length 3, given by the two first and the last letters of $\mathbf{w}$ (they are circled points in Figure 11, right). By Theorem 1.11, $\mathbf{w}$ is not CFC, unless $s_i = u$, and therefore $\mathbf{w}$ satisfies (c).

If $s_j = s_{i+1}$, then the last but one letter in $\mathbf{w}$ is either s_i or s_{i+2} , so H^c takes one of the two forms in Figure 12.




FIGURE 12. The two possible heaps for $s_j = s_{i+1}$.

In the first case, H^c contains a cylindric convex chain involving points in $H_{\{s_i, s_{i+1}\}}^c$, of length 3, given by the first and the two last letters of $\mathbf{w}$ (the corresponding points are circled in Figure 12, left). By Theorem 1.11, $\mathbf{w}$ is not CFC, unless $s_i = s_0 = t$, which is not possible because the second letter of $\mathbf{w}$ is s_{i-1} , or unless $s_{i+1} = s_n = u$, and therefore $\mathbf{w}$ satisfies (c). In the second case, $\mathbf{w}$ satisfies (c).

- w belongs to one of the three remaining families, called right peak, left peak and left-right peak. In this case, it is easy to find either a chain covering relation $p \prec_c q$ such that p and q have the same label or a cylindric convex chain in H^c , except if w satisfies (c) (and has length $2n$).

Conversely, any w satisfying one of the three conditions (a), (b), or (c) is a CFC element. Indeed, H^c contains neither a relation $i \prec_c j$ such that i and j have the same label nor a cylindric convex chain of length 3, except involving points in $H_{\{t, s_1\}}^c$ or $H_{\{s_{n-1}, u\}}^c$ which do not contain any cylindric convex chain of length 4: for w satisfying (a) or (c), this is trivial, and for w satisfying (b), the same proof as in type $\tilde{A}$ holds (see Theorem 2.1). $\square$

Corollary 3.2. *We have the generating function*

$$\tilde{C}_n^{CFC}(q) = A_{n+1}^{CFC}(q) + \frac{2^n}{1 - q^{n+1}} q^{2(n+1)} + \frac{2n}{1 - q^{2n}} q^{2n}. \quad (5)$$

The coefficients of $\tilde{C}_n^{CFC}(q)$ are ultimately periodic of exact period $n(n+1)$ if n is odd, and $2n(n+1)$ if n is even. Moreover, periodicity starts at length n .

Proof. Notice that the sets of elements satisfying condition (a), (b) or (c) of Theorem 3.1 are disjoint. The first term in (5) corresponds to the set of CFC elements satisfying (a). This set is clearly in bijection with A_{n+1}^{CFC} .

Now, we show that the second term in (5) corresponds to CFC elements satisfying (b). If we fix an integer $h \geq 2$, there are 2^n such elements satisfying $h = |\mathbf{w}_t| = |\mathbf{w}_{s_1}| = \dots = |\mathbf{w}_{s_{n-1}}| = |\mathbf{w}_u|$. Indeed, for each generator $t, s_1, \dots, s_{n-1}$, there are two possibilities: s_i appears before or after s_{i+1} (and t appears before or after s_1). The generating function of elements satisfying (b) is then equal to $\sum_{h \geq 2} 2^n q^{h(n+1)}$.

With the condition on the number of occurrences of each generator, we can see that the length of elements satisfying (c) must be a multiple of $2n$, and there are $2n$ such elements for each possible length (the first two letters in $\mathbf{w}$ can be $s_i s_{i+1}$ or $s_i s_{i-1}$ for all $i \in \llbracket 1, n-1 \rrbracket$, or ts_1 or us_{n-1}). This gives the term $\frac{2n}{1-q^{2n}} q^{2n}$.

$A_{n+1}^{CFC}(q)$ is a polynomial in q , and the two other terms in (5) have periodic coefficients, so $\tilde{C}_n^{CFC}(q)$ is ultimately periodic. The period is the least common multiple of $n+1$ and $2n$, which is $n(n+1)$ if n is odd, and $2n(n+1)$ if n is even. As the leading coefficient of $A_{n+1}^{CFC}(q)$ is $2^n q^{n+1}$ and $[q^n]A_{n+1}^{CFC}(q) \neq 0$, the beginning of the periodicity is n . $\square$

We can deduce the characterization in type B , whose corresponding Coxeter diagram is recalled below.




FIGURE 13. Coxeter diagram of type B_n .


Corollary 3.3. *The CFC elements in type B_n are those having reduced expressions in which each generator occurs at most once. Moreover,*

$$B_n^{CFC}(q) = A_n^{CFC}(q).$$

Proof. Let w be a CFC element in type B_n , and $\mathbf{w}$ one of its reduced expression. Consequently, w is a CFC element in type $\tilde{C}_n$ in which the generator u does not occur. But, according to Theorem 2.1, the only such CFC elements are those satisfying (a) (and having no u): if w satisfies (b) (*resp.* (c)), all generators appear the same number of times (*resp.* u must appear in $\mathbf{w}$). Conversely, if all generators occur at most once in $\mathbf{w}$, we already saw in Remark 1.13 that w is a CFC element. $\square$

3.2. Types $\tilde{B}_{n+1}$ and D_{n+1} . We will obtain once again a characterization of CFC elements in type $\tilde{B}_{n+1}$, deduce a characterization and the enumeration in type D_{n+1} , and finally deduce the enumeration in type $\tilde{B}_{n+1}$.

In what follows, by “ $\mathbf{w}$ is an alternating word”, we mean that if we replace t_1 and t_2 by s_0 , and u by s_n , $\mathbf{w}$ is an alternating word in the sense of Definition 1.12, and t_1 and t_2 alternate in $\mathbf{w}$.

FIGURE 14. Coxeter diagrams of types $\tilde{B}_{n+1}$ and D_{n+1} .

Theorem 3.4. *An element $w \in \tilde{B}_{n+1}$ is CFC if and only if one (equivalently any) of its reduced expressions $\mathbf{w}$ verifies one of these conditions:*

- (a) *each generator occurs at most once in $\mathbf{w}$,*
- (b) *$\mathbf{w}$ is an alternating word, $|\mathbf{w}_{s_1}| = \dots = |\mathbf{w}_{s_{n-1}}| = |\mathbf{w}_u| \geq 2$, and $|\mathbf{w}_{t_1}| = |\mathbf{w}_{t_2}| = |\mathbf{w}_{s_1}|/2$ (in particular, $|\mathbf{w}_{s_1}|$ is even),*
- (c) *$\mathbf{w}$ is a subword of $(t_1 t_2 s_1 s_2 \dots s_{n-1} u s_{n-1} \dots s_2 s_1)^\infty$, which is an infinite periodic word, where t_1 and t_2 are allowed to commute, such that $|\mathbf{w}_{s_1}| = \dots = |\mathbf{w}_{s_{n-1}}| \geq 2$, and $|\mathbf{w}_{t_2}| = |\mathbf{w}_{t_1}| = |\mathbf{w}_u| = |\mathbf{w}_{s_1}|/2$ (i.e $\mathbf{w}$ takes one of the five forms: $s_i s_{i+1} \dots s_{i-2} s_{i-1}$ or $s_i s_{i-1} \dots s_{i+2} s_{i+1}$ or $t_1 s_1 \dots s_1 t_2$ or $t_2 s_1 \dots t_1$ or $t_1 t_2 s_1 \dots s_1$).*

Proof. The steps of this proof are the same as in Theorem 3.1. Let w be a CFC element in $\tilde{B}_{n+1}$ and let $\mathbf{w}$ be one of its reduced expressions. We denote by H the heap of $\mathbf{w}$ and H^c its cylindric transformation. In [1, Theorem 3.10], FC elements are classified in five families. The first corresponds to alternating elements. As before, we distinguish two cases:

- each generator occurs at most once in $\mathbf{w}$. These elements satisfy (a).
- $\mathbf{w}$ comes from an alternating word of type $\tilde{C}_n$, where we applied the replacements $(t, t, \dots, t) \rightarrow (t_1, t_2, t_1, \dots)$ or $(t_2, t_1, \dots)$. The same proof as for type $\tilde{C}$ gives $|\mathbf{w}_t| = |\mathbf{w}_{s_1}| = \dots = |\mathbf{w}_{s_{n-1}}| = |\mathbf{w}_u|$. If $|\mathbf{w}_t|$ is odd, after replacement we have the subword of $\mathbf{w}$ where we keep only the generators t_1, t_2 and s is $t_1 s_1 t_2 s_1 \dots s_1 t_1$ (or $t_2 s_1 t_1 s_1 \dots s_1 t_2$), which prevents w from being CFC. So $|\mathbf{w}_t|$ is even, and as $\mathbf{w}$ is alternating, $|\mathbf{w}_{t_2}| = |\mathbf{w}_{t_1}| = |\mathbf{w}_{s_1}|/2$. These elements satisfy (b).

Next, we will show that among the four remaining families from [1], the CFC elements must satisfy (c). There are two possibilities:

- $\mathbf{w}$ is a subword of $(t_1 t_2 s_1 s_2 \dots s_{n-1} u s_{n-1} \dots s_2 s_1)^\infty$, which is an infinite periodic word, where t_1 and t_2 are allowed to commute, such that a generator occurs more than twice. The same case distinction as in type $\tilde{C}$ leads us to the required condition (c).
- $H_{\mathbf{w}}$ is a heap among special cases analogous to the ones in type $\tilde{C}_n$ which are trivially non CFC, except for those which satisfy (c) (and are of length $2(n+2)$).

Conversely, all elements w satisfying one of the three conditions (a), (b) or (c) are CFC elements. Indeed, H^c contains neither relations $i \prec j$ such that i and j have the same label nor cylindric convex chains of length 3, except involving points in $H_{\{s_{n-1}, u\}}^c$, and these points can not form a cylindric convex chain of length 4. $\square$

Proposition 3.5. *The CFC elements in type D_{n+1} are those having reduced expressions in which each generator occurs at most once. Moreover, we have $D_1^{CFC}(q) =$*

$1 + q$, $D_2^{CFC}(q) = 1 + 2q + q^2$, $D_3^{CFC}(q) = 1 + 3q + 5q^2 + 4q^3$ and for $n \geq 3$:

$$D_{n+1}^{CFC}(q) = (2q + 1)D_n^{CFC}(q) - qD_{n-1}^{CFC}(q). \quad (6)$$

In other words, we have

$$D^{CFC}(x) := \sum_{n=0}^{\infty} D_{n+1}^{CFC}(q)x^n = \frac{(1+q) - xq(1+q) + x^2q^2(1+2q)}{1 - (2q+1)x + qx^2}.$$

Proof. Let w be a CFC element of type D_{n+1} and let $\mathbf{w}$ be one of its reduced expressions. Consequently, w is a CFC element of type $\tilde{B}_{n+1}$ in which the generator u does not occur. But, according to Theorem 3.4, the only such CFC elements are those satisfying condition (a) (and having no u): if $\mathbf{w}$ satisfies (b) or (c), u must appear in $\mathbf{w}$. Conversely, if all generators occur at most once in $\mathbf{w}$, we already saw that w is CFC.

So, any CFC element of type D_{n+1} can be obtained from a CFC element w' in A_{n-1} by adding to w' nothing, or one occurrence of t_1 (with two choices: before or after s_1), or one occurrence of t_2 (with two choices), or one occurrence of t_1 and t_2 (with four choices). If s_1 does not occur in w' , we have no choice for adding t_1 or/and t_2 as t_1 and t_2 commute with all other generators. This leads to the following recurrence relation:

$$D_{n+1}^{CFC}(q) = (1 + 4q + 4q^2)A_{n-1}^{CFC}(q) - (2q + 3q^2)A_{n-2}^{CFC}(q).$$

With this relation and (1) in Theorem 2.3, it is easy to derive (6). The generating function $D^{CFC}(x)$ is computed through classical methods. $\square$

Corollary 3.6. *We have the generating function*

$$\tilde{B}_{n+1}^{CFC}(q) = D_{n+2}^{CFC}(q) + \frac{2^{n+1}}{1 - q^{2(n+1)}} q^{2(n+1)} + \frac{2(n+1)}{1 - q^{2n+1}} q^{2n+1}. \quad (7)$$

Furthermore, the coefficients of $\tilde{B}_{n+1}^{CFC}(q)$ are ultimately periodic of exact period $2(n+1)(2n+1)$ and the periodicity starts at length $n+2$.


Proof. Notice that the sets of elements satisfying condition (a), (b) or (c) of Theorem 3.4 are disjoint. The first term in (7) corresponds to the set of CFC elements satisfying (a). This set is clearly in bijection with D_{n+2}^{CFC} .

The second term in (7) corresponds to CFC elements satisfying (b), which have length a multiple of $2(n+1)$, and there are 2^{n+1} such elements for each possible length.

With the condition on the number of occurrences of each generator, we can see that the length of elements satisfying (c) must be a multiple of $2n+1$, and there are $2n+2$ such elements for each possible length (the first two letters can be $s_i s_{i+1}$, $s_i s_{i-1}$, $u s_{n-1}$, $t_1 t_2$, $t_1 s_1$ or $t_2 s_1$). This gives the third term of (7).

Moreover, $D_{n+2}^{CFC}(q)$ is a polynomial in q , and the two other terms of (7) are periodic, so $\tilde{B}_{n+1}^{CFC}(q)$ is ultimately periodic. The period is the least common multiple of $2(n+1)$ and $2n+1$, which is $2(n+1)(2n+1)$. The beginning of the periodicity is $n+2$, as the leading coefficient of $D_{n+2}^{CFC}(q)$ is $2^{n+1}q^{n+2}$ and $[q^{n+1}]D_{n+2}^{CFC}(q) \neq 0$. $\square$

3.3. Type $\tilde{D}_{n+2}$. The situation is very similar in type $\tilde{D}_{n+2}$ for the characterization, but the generating function takes a slightly different form, due to the specificity of the Coxeter diagram.

FIGURE 15. Coxeter diagram of type $\tilde{D}_{n+2}$.

In what follows, by “ $\mathbf{w}$ is an alternating word”, we mean that if we replace t_1 and t_2 by s_0 , and u_1 and u_2 by s_n , the image of $\mathbf{w}$ is an alternating word in the sense of Definition 1.12, t_1 and t_2 alternate in w , and u_1 and u_2 alternate in w .

Theorem 3.7. *An element $w \in \tilde{D}_{n+2}$ is CFC if and only if one (equivalently any) of its reduced expressions $\mathbf{w}$ verifies one of these conditions:*

- (a) *each generator occurs at most once in $\mathbf{w}$,*
- (b) *$\mathbf{w}$ is an alternating word, $|\mathbf{w}_{s_1}| = \dots = |\mathbf{w}_{s_{n-1}}| \geq 2$, and $|\mathbf{w}_{t_1}| = |\mathbf{w}_{t_2}| = |\mathbf{w}_{u_1}| = |\mathbf{w}_{u_2}| = |\mathbf{w}_{s_1}|/2$ (in particular, $|\mathbf{w}_{s_1}|$ is even),*
- (c) *$\mathbf{w}$ is a subword of $(t_1 t_2 s_1 s_2 \dots s_{n-1} u_1 u_2 s_{n-1} \dots s_2 s_1)^\infty$, which is an infinite periodic word, where t_1 and t_2 , u_1 and u_2 are allowed to commute, such that $|\mathbf{w}_{s_1}| = \dots = |\mathbf{w}_{s_{n-1}}| \geq 2$, and $|\mathbf{w}_{t_2}| = |\mathbf{w}_{t_1}| = |\mathbf{w}_{u_1}| = |\mathbf{w}_{u_2}| = |\mathbf{w}_{s_1}|/2$.*

Proof. The same proof as for Theorem 3.4 holds, one only needs to add the replacements $(u, u, \dots u) \rightarrow (u_1, u_2, \dots)$ or $(u_2, u_1, \dots)$. $\square$

Again, we can compute the corresponding generating function.

Proposition 3.8. *We have the generating function*

$$\tilde{D}_{n+2}^{CFC}(q) = Q_{n+2}(q) + \frac{2^{n+2} + 2(n+2)}{1 - q^{2(n+1)}} q^{2(n+1)}, \quad (8)$$

where $Q_{n+2}(q)$ is a polynomial in q of degree $n+3$ such that $Q_4(q) = 1 + 5q + 14q^2 + 28q^3 + 33q^4 + 16q^5$, $Q_5(q) = 1 + 6q + 20q^2 + 46q^3 + 73q^4 + 72q^5 + 32q^6$, and for $n \geq 4$:

$$Q_{n+2}(q) = (2q + 1)Q_{n+1}(q) - qQ_n(q).$$

Moreover, the coefficients of $\tilde{D}_{n+2}^{CFC}(q)$ are ultimately periodic of exact period $2(n+1)$, and the periodicity starts at length $n+2$.

Note that the generating function $Q(x)$ of the polynomials $Q_n(q)$ is computable through classical techniques. However it does not have a nice expression.

Proof. Notice that the sets of elements satisfying condition (a), (b) or (c) of Theorem 3.7 are disjoint. The first term in (8) corresponds to the set of CFC elements satisfying (a). Any such element can be obtained from a CFC element w of type D_{n+1} by adding nothing, or one occurrence of u_1 (with two choices: before or after s_{n-1}), or one occurrence of u_2 (with two choices), or one occurrence of u_1 and u_2 (with four choices). If s_{n-1} does not occur in w , we have no choice for adding u_1 or/and u_2 , as u_1 and u_2 commute with all other generators. This leads to the following recurrence relation:

$$Q_{n+2}(q) = (1 + 4q + 4q^2)D_{n+1}^{CFC}(q) - (2q + 3q^2)D_n^{CFC}(q).$$

Using this and (6), we find the expected recurrence relation for the polynomials $Q_n(q)$. Elements satisfying (b) have length a multiple of $2(n+1)$ and there are

or an integer multiple of a constant depending on the considered type.



FIGURE 16. Coxeter diagrams of types $\tilde{G}_2$, $\tilde{E}_6$ and $\tilde{E}_7$.

(equivalently any) of its reduced expressions $\mathbf{w}$ verifies one of these conditions:

- W₂.

Consequently, the following generating functions hold:

$$\tilde{G}_2^{CFC}(q) = R_1(q) + \frac{6q^5}{1 - q^5},$$

$$\tilde{E}_6^{CFC}(q) = R_2(q) + \frac{23q^{12}}{1 - q^{12}},$$

$$\tilde{E}_7^{CFC}(q) = R_3(q) + \frac{45q^{18}}{1-q^{18}},$$

where $R_1(q)$, $R_2(q)$ and $R_3(q)$ are polynomials.

3.5. Logarithmic CFC elements. Here we study some particular elements of a general Coxeter group W . Recall that for $w \in W^{CFC}$, the support $\text{supp}(w)$ of w is the set of generators that appear in a (equivalently, any) reduced expression of w . It is well known that if $w \in W$ and k is a positive integer, then $\ell(w^k) \leq k\ell(w)$. If equality holds for all $k \in \mathbb{N}^*$ (i.e. w^k is reduced for all k), we say that w is logarithmic (see [3] for more information about logarithmic elements). For example, it is trivial that there is no logarithmic element in finite Coxeter groups, because there is a finite number of reduced expressions. By using our characterizations for affine types in Theorems 2.1, 3.4, 3.1, 3.7, we derive the following consequence. A generalization of this result was proved for all Coxeter groups in [8, Corollary E], by using geometric group theoretic methods. Nevertheless, as our approach is different and more combinatorial, we find interesting to give our proof, although it only works for affine types.

Theorem 3.10. *For $W = \tilde{A}, \tilde{B}, \tilde{C}$, or $\tilde{D}$, if w is a CFC element, w is logarithmic if and only if a (equivalently, any) reduced expression $\mathbf{w}$ of w has full support (i.e. all generators occur in $\mathbf{w}$). In particular, there is a finite number of CFC elements which are not logarithmic.*

Proof. Let w be in W^{CFC} and let $\mathbf{w}$ be one of its reduced expressions. Assume that $\mathbf{w}$ has not full support. In this case, w belongs to a proper parabolic subgroup of W , which is a finite Coxeter group by a classical property of affine Coxeter groups (see [7]). So w is not logarithmic, and there is a finite number of such elements, which correspond to elements satisfying (a) in Theorems 2.1, 3.4, 3.1, 3.7 and such that at least one generator does not occur in $\mathbf{w}$.

Conversely, assume that w is a CFC element with full support. According to Theorems 2.1, 3.4, 3.1, 3.7, $\mathbf{w}$ must satisfy (b) or (c) or has to be a Coxeter element (i.e. each generator occurs exactly once in $\mathbf{w}$). If w is a Coxeter element, w is logarithmic (see [14]). If $\mathbf{w}$ satisfies (b) (resp. (c)), it is easy to verify that $\mathbf{w}^k$ also satisfies (b) (resp. (c)) and is therefore reduced. It follows that w is logarithmic. $\square$

We can also notice that the powers of Coxeter elements are always CFC in affine types $\tilde{A}_{n-1}$ and $\tilde{C}_n$ because they satisfy the alternating word condition, but are never CFC elements in types $\tilde{B}_{n+1}$ and $\tilde{D}_{n+2}$ (because they satisfy neither the alternating word condition nor condition (c)).

4. CFC INVOLUTIONS

A natural question that arises in the study of FC elements is to compute the number of FC involutions in finite and affine Coxeter groups (as this number is, for some groups, the sum of the dimensions of irreducible representations of a natural quotient of the Iwahori-Hecke algebra associated to the group, see [5]). Similarly, we now focus on CFC elements which are involutions. The main result is that there is always a finite number of such elements in all Coxeter groups, and we are able to characterize them in terms of words. We also use the characterization of CFC elements to enumerate CFC involutions in finite and affine Coxeter groups.

4.1. Finiteness and characterization of CFC involutions.

Theorem 4.1. *Let W be a Coxeter group and let $I(W)$ be its subset of involutions. The set $W^{CFC} \cap I(W)$ is finite. Moreover an element w belongs to $W^{CFC} \cap I(W)$ if*

and only if for any generator s in $\text{supp}(w)$, $|w_s| = 1$ and for all t such that $m_{st} \geq 3$, $|w_t| = 0$ (i.e two non commuting generators can not occur in w).

Proof. Let w be a CFC involution, let s be a generator in $\text{supp}(w)$ and $\mathbf{w}$ be a reduced expression of w . Assume that $|w_s| \geq 2$. Consider a cyclic shift $\mathbf{w}' = s\mathbf{w}_1$ of $\mathbf{w}$ which begins with s . As w and s are involutions, $\mathbf{w}'$ is an expression of an involution. Moreover as w is CFC, $\mathbf{w}'$ corresponds to a FC element w' . According to [11], a FC element w is an involution if and only if $R(w)$ is palindromic, i.e $R(w)$ includes the reverse of all of its members. Applying this to $\mathbf{w}'$ allows us to say that $\mathbf{w}'$ is commutation equivalent to a word $s\mathbf{w}_2s$. So a cyclic shift of $\mathbf{w}$ is commutation equivalent to $s\mathbf{w}_2s$, which is in contradiction with the CFC property of w . Therefore we have $|w_s| = 1$. We consider $\mathbf{w}'$ as before. As $R(w')$ is palindromic, we can conclude that all generators in $\text{supp}(w') = \text{supp}(w)$ commute with s . $\square$

Remark 4.2. The number of CFC involutions in a Coxeter group W depends only on the edges of the Coxeter diagram, without taking into account the values m_{st} .

4.2. CFC involutions in classical types. Let us enumerate the CFC involutions in classical types, according to their Coxeter length. If W is a Coxeter group, we define $W^{CFCI}(q) := \sum_{w \in W^{CFC} \cap I(W)} q^{\ell(w)}$.

Theorem 4.3. *In types A , B , and $\tilde{C}$ the following relation holds for all $n \geq 2$:*

$$A_n^{CFCI}(q) = B_n^{CFCI}(q) = \tilde{C}_{n-1}^{CFCI}(q) = A_{n-1}^{CFCI}(q) + qA_{n-2}^{CFCI}(q), \quad (9)$$

and $A_0^{CFCI}(q) = 1$, $A_1^{CFCI}(q) = 1 + q$. Moreover, we can compute the generating function

$$A^{CFCI}(x) := \sum_{n=1}^{\infty} A_{n-1}^{CFCI}(q)x^n = x \frac{1+qx}{1-x-qx^2}.$$

Proof. The equality $A_n^{CFCI}(q) = B_n^{CFCI}(q) = \tilde{C}_{n-1}^{CFCI}(q)$ comes from Remark 4.2. Let w be a CFC involution in A_n . If s_n belongs to $\text{supp}(w)$, by Theorem 4.1, s_{n-1} does not belong to $\text{supp}(w)$, and w is equal to s_n concatenated to a CFC involution of A_{n-2} . If s_n does not belong to $\text{supp}(w)$, w is a CFC involution of A_{n-1} . This yields the expected relation (9). The generating function is computed through classical techniques. $\square$

In particular, if $q \rightarrow 1$, we obtain the number of CFC involutions in type A_{n-1} , which is the $(n+1)^{\text{th}}$ Fibonacci number.

Theorem 4.4. *In types D and $\tilde{B}$, the following relations hold for all $n \geq 3$:*

$$D_n^{CFCI}(q) = \tilde{B}_{n-1}^{CFCI}(q) = D_{n-1}^{CFCI}(q) + qD_{n-2}^{CFCI}(q), \quad (10)$$

$$D_{n+1}^{CFCI}(q) = qA_{n-3}^{CFCI}(q) + (1+2q+q^2)A_{n-2}^{CFCI}(q), \quad (11)$$

and $D_1^{CFCI}(q) := 1$, $D_2^{CFCI}(q) = 1+2q+q^2$, $D_3^{CFCI}(q) = 1+3q+q^2$. Moreover, we can compute the generating function

$$D^{CFCI}(x) := \sum_{n=0}^{\infty} D_{n+1}^{CFCI}(q)x^n = \frac{1+(2q+q^2)x}{1-x-qx^2}.$$

Proof. To prove (11), let w be a CFC involution in D_{n+1} . If s_1 belongs to $\text{supp}(w)$, by Theorem 4.1, s_2, t_1, t_2 do not belong to $\text{supp}(w)$, and w is equal to s_1 concatenated to a CFC involution of A_{n-3} . If s_1 does not belong to $\text{supp}(w)$, w is a CFC involution of A_{n-1} concatenated to t_1, t_2, t_1t_2 or nothing. This shows (11). Equation (10) is simply a consequence of (11), (9) and Remark 4.2. The generating function comes from classical techniques. $\square$

If $q \rightarrow 1$, we obtain the number of CFC involutions in type D_{n+1} , which leads to a Fibonacci-type sequence with starting numbers 1 and 4.

Theorem 4.5. *In type $\tilde{A}$, the following relations hold for $n \geq 2$:*

$$\tilde{A}_n^{CFCI}(q) = \tilde{A}_{n-1}^{CFCI}(q) + q\tilde{A}_{n-2}^{CFCI}(q), \quad (12)$$

where $\tilde{A}_0^{CFCI}(q) := 1$, $\tilde{A}_1^{CFCI}(q) = 1 + 2q$. Therefore, we can compute the generating function

$$\tilde{A}^{CFCI}(x) := \sum_{n=1}^{\infty} \tilde{A}_{n-1}^{CFCI}(q)x^n = x \frac{1 + 2qx}{1 - x - qx^2}.$$

Remark 4.6. We also have

$$q^n \tilde{A}_{n-1}^{CFCI}(1/q^2) = L_n(q),$$

where $L_n(q)$ is the n^{th} Lucas polynomials, defined explicitly (see sequence A114525 in [10]) by $L_n(q) = 2^{-n}[(q - \sqrt{q^2 + 4})^n + (q + \sqrt{q^2 + 4})^n]$. This equality can be proved by using generating functions.

Proof. Let w be a CFC involution in $\tilde{A}_{n-1}$. If s_0 belongs to $\text{supp}(w)$, by Theorem 4.1, s_{n-1} and s_1 do not belong to $\text{supp}(w)$, and w is equal to s_0 concatenated to a CFC involution of A_{n-3} . If s_0 does not belong to $\text{supp}(w)$, w is a CFC involution of A_{n-1} . This leads to the relation:

$$\tilde{A}_{n-1}^{CFCI}(q) = qA_{n-3}^{CFCI}(q) + A_{n-1}^{CFCI}(q).$$

Using this and (9), it is easy to deduce (12). The generating function comes from classical techniques. $\square$

Theorem 4.7. *In type $\tilde{D}$, the following relation holds for $n \geq 4$:*

$$\tilde{D}_{n+2}^{CFCI}(q) = \tilde{D}_{n+1}^{CFCI}(q) + q\tilde{D}_n^{CFCI}(q), \quad (13)$$

with $\tilde{D}_4^{CFCI}(q) = 1 + 5q + 6q^2 + 4q^3 + q^4$ and $\tilde{D}_5^{CFCI}(q) = 1 + 6q + 10q^2 + 6q^3 + q^4$. We can therefore compute the generating function:

$$\tilde{D}^{CFCI}(x) := \sum_{n=2}^{\infty} \tilde{D}_{n+2}^{CFCI}(q)x^n = x^2 \frac{1 + 5q + 6q^2 + 4q^3 + q^4 + x(q + 4q^2 + 2q^3)}{1 - x - qx^2}.$$

Proof. Let w be a CFC involution in $\tilde{D}_{n+2}$. If s_1 belongs to $\text{supp}(w)$, by Theorem 4.1, s_2, t_1 and t_2 do not belong to $\text{supp}(w)$, and w is equal to s_1 concatenated to a CFC involution of D_{n-1} . If s_1 does not belong to $\text{supp}(w)$, w is a CFC involution D_n concatenated to t_1, t_2, t_1t_2 or nothing. This leads to the relation:

$$\tilde{D}_{n+2}^{CFCI}(q) = qD_{n-1}^{CFCI}(q) + (1 + 2q + q^2)D_n^{CFCI}(q).$$

Use this and (10) to derive (13). $\square$

If $q \rightarrow 1$, we obtain the number of involutions in type $\tilde{D}_{n+2}$, which is a Fibonacci-type sequence with starting numbers 10 and 7.

5. OTHER QUESTIONS

In Sections 2 and 3, we obtained two q -analogs of the number of CFC elements in finite types. The first, in type A , corresponds to permutations avoiding 321 and 3412, taking into account their Coxeter length (see [15]). The second, in type D , is apparently new. One may wonder if it corresponds to another combinatorial object. We can notice that if W is a finite or affine Coxeter group, the generating function of CFC elements is rational. This is in fact true for all Coxeter groups, as will be proved in the forthcoming work [9].

6. ACKNOWLEDGEMENTS

We would like to thank M. Macauley for pointing out reference [8] on logarithmic elements, and T. Marquis for valuable remarks on the infinite number of non logarithmic CFC elements. Finally, we also thank P. Nadeau for interesting comments and discussions on an earlier version of this manuscript.

REFERENCES

- [1] R. Biagioli, F. Jouhet, and P. Nadeau, *Fully commutative elements in finite and affine Coxeter groups*, to appear in Monatsh. Math., preprint arXiv:1402.2166 (2014), 37 pages.
- [2] A. Björner, and F. Brenti, *Combinatorics of Coxeter groups*, Springer (2005).
- [3] T. Boothby, J. Burkert, M. Eichwald, D. C. Ernst, R. M. Green, and M. Macauley, *On the cyclically fully commutative elements of Coxeter groups*, J. Algebraic Combin. **1** (2012), 123–148.
- [4] C. K. Fan, *A Hecke algebra quotient and properties of commutative elements of a Weyl group.*, Phd thesis, M.I.T. (1995).
- [5] C. K. Fan, *Structure of a Hecke Algebra Quotient*, J. Amer. Math. Soc. **1**(1997), 139–167.
- [6] J. Graham, *Modular Representations of Hecke Algebras and Related Algebras*, PhD thesis, University of Sydney (1995).
- [7] J. E. Humphreys, *Reflection groups and Coxeter groups*, Volume 29 of Cambridge Studies in Advanced Mathematics, Cambridge University Press, Cambridge (1990).
- [8] T. Marquis, *Conjugacy classes and straight elements in Coxeter groups*, to appear in Journal of Algebra, preprint arXiv:1310.1021v2 (2014), 12 pages.
- [9] M. Pétreolle, *Rational generating function for cyclically fully commutative elements*, in preparation (2014).
- [10] N. J. A. Sloane, *The on-line encyclopedia of integer sequences* (2013).
- [11] J. R. Stembridge, *On the fully commutative elements of Coxeter groups*, J. Algebraic Combin. **4** (1996), 353–385.
- [12] J. R. Stembridge, *Some combinatorial aspects of reduced words in finite Coxeter groups*, Trans. Amer. Math. Soc. **4** (1997), 1285–1332.
- [13] J. R. Stembridge, *The enumeration of fully commutative elements of Coxeter groups*, J. Algebraic Combin. **7** (1998), 291–320.
- [14] D. E. Speyer, *Powers of Coxeter elements in infinite groups are reduced* Proc. Amer. Math. Soc. **4** (2009), 1295–1302.
- [15] B. E. Tenner, *Pattern avoidance and the Bruhat order*, J. Combin. Theory Ser. A **5** (2007), 888–905.

INSTITUT CAMILLE JORDAN, UNIVERSITÉ CLAUDE BERNARD LYON 1, 69622 VILLEURBANNE CEDEX, FRANCE

E-mail address: petreolle@math.univ-lyon1.fr

URL: <http://math.univ-lyon1.fr/~petreolle>